

Metatarsalgia - forefoot pain

What is metatarsalgia?

This is a foot condition, in which discomfort is felt under the forefoot region, often referred to as the “ball of the foot”. This is a generic name, which may have many underlying causes.



What are the symptoms?

The pain is varied and can range from mild to severe discomfort and is often made worse with standing or walking. It is sometimes described as a burning or aching sensation, sometimes sharp or you may have shooting pains, tingling or numbness in your toes. Some people also experience a sensation that feels like walking on pebbles. The symptoms of metatarsalgia usually develop gradually and can get worse over time.

What can cause this condition?

There are many causes to metatarsalgia. These may include:

- Certain footwear may predispose you to this condition, including high heels, tight fitting shoes and shoes with little cushioning.
- Being overweight or obese - this can alter your foot function and increase stresses through the foot
- Foot type and function - this may include a “flatter” or “high arch foot type”
- Foot deformities, such as hammer toes, may predispose you to pain in the forefoot
- Age - the fat pad that protects the foot can thin with age, making patients more likely to feel pain in the ball of their foot
- Tight muscles such as the calf muscle, which can cause increased pressures through the front of the foot
- Stress fractures in the foot - these occasionally occur in athletes or walkers and cause pain to come on rapidly, often with some degree of swelling.
- Inflamed small nerves known as neuromas, can cause forefoot pain.

Some medical conditions may lead to pain in the “balls of the feet”. These may include:

- Inflammatory arthritis, such as rheumatoid arthritis or gout
- Osteoarthritis - A “wear and tear” type arthritis
- Diabetes - which can lead to nerve damage and an altered foot shape

How is my condition diagnosed?

A foot care practitioner may diagnose your condition by examining the foot. In some cases blood tests and imaging such as ultrasound may be required.

How is my condition treated?

Treatments will depend on the underlying cause of your condition. In most cases of metatarsalgia



Metatarsalgia - forefoot pain

you will need to ensure that:

1. Calf stretches are carried out regularly - as directed by your practitioner
2. Appropriate footwear is worn, with cushioning and adequate width. Avoid footwear such as high heels which direct more pressure under the forefoot regions.
3. Orthoses (medical shoe insoles) may be prescribed to reduce the pressure under the forefoot and improve foot function
4. Weight-loss is important if you are overweight.
5. Pain Killers - may help to reduce some of your symptoms

Other specialist interventions:

Injections may be useful for some cases of metatarsalgia

Surgery may be required in a number of cases, (such as toe straightening) but this will depend on the underlying cause and your suitability for surgery

Prescribed medication may be given if the cause of the foot pain is due to a systemic condition such as rheumatoid arthritis.